



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C12N 15/54, 15/82, 9/10, C12Q 1/02, 1/68, 1/48</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/37656</b> <b>(43) International Publication Date:</b> 29 June 2000 (29.06.00)
<b>(21) International Application Number:</b> PCT/US99/30338 <b>(22) International Filing Date:</b> 20 December 1999 (20.12.99) <b>(30) Priority Data:</b> 60/113,190 21 December 1998 (21.12.98) US <b>(71) Applicant (for all designated States except US):</b> E.I. DU PONT DE NEMOURS AND COMPANY [US/US]; 1007 Market Street, Wilmington, DE 19898 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> FAMODU, Omolayo, O. [US/US]; 216 Barrett Run Place, Newark, DE 19702 (US). MCGONIGLE, Brian [US/US]; 1707 North Union Street, Wilmington, DE 19806 (US). ODELL, Joan, T. [US/US]; P.O. Box 826, Unionville, PA 19375 (US). FADER, Gary, M. [US/US]; 1000 Woods Lane, Landenberg, PA 19350 (US). FALCO, Saverio, Carl [US/US]; 1902 Miller Road, Arden, DE 19810 (US). <b>(74) Agent:</b> FEULNER, Gregory, J.; E.I. du Pont de Nemours and Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).		<b>(81) Designated States:</b> AE, AL, AU, BA, BB, BG, BR, CA, CN, CR, CU, CZ, DM, EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> FLAVONOID BIOSYNTHETIC ENZYMES  <b>(57) Abstract</b>  This invention relates to an isolated nucleic acid fragment encoding a flavonoid biosynthetic enzyme. The invention also relates to the construction of a chimeric gene encoding all or a portion of the flavonoid biosynthetic enzyme, in sense or antisense orientation, wherein expression of the chimeric gene results in production of altered levels of the flavonoid biosynthetic enzyme in a transformed host cell.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TITLE

## FLAVONOID BIOSYNTHETIC ENZYMES

This application claims the benefit of U.S. Provisional Application No. 60/113,190, filed December 21, 1998.

FIELD OF THE INVENTION

This invention is in the field of plant molecular biology. More specifically, this invention pertains to nucleic acid fragments encoding flavonoid biosynthetic enzymes in plants and seeds.

BACKGROUND OF THE INVENTION

Plants accumulate a variety of natural products that are synthesized in response to environmental stimuli and genetically programmed developmental signals. Of these secondary metabolites, flavonoids are probably the most ubiquitous. Flavonoids are products of biosynthetic pathways originating from the central phenylpropanoid pathway and the acetate-malonate pathway. Flavonoids represent a large class of molecules that serve many diverse functions, such as co-pigments in flower color, stimulation of pollen tube growth, pollinator attraction, and feeding deterrents and protection against UV irradiation in fruits and seeds (Holton, T. A. et al. (1993) *Plant J* 4:1003-1010). Therefore, there has been considerable interest in understanding flavonoid biosynthesis, which in turn would afford the ability to control pigmentation and many other phenotypic traits associated with flavonoid synthesis.

Isoflavonoids constitute a large group of naturally occurring flavonoids. They occur almost exclusively in leguminous plants and form part of the host defense response against phytopathogenic microorganisms (Akashi, T. et al. (1998) *Biochemical and Biophysical Research Communications* 251:67-70). For example, a variety of simple isoflavonoids like coumestans, pterocarpans and isoflavans have been shown to have antimicrobial properties.

An essential part of the biosynthesis of these compounds is the 2'-hydroxylation of isoflavones by isoflavone 2'-hydroxylase (I2'H). It may be possible to change the levels of some isoflavones in plants by altering the level or activity of isoflavone biosynthetic enzymes. One unique isoflavone, daidzein (5-deoxyisoflavonoid), is a product of the phenylpropanoid pathway and is a healthful compound found in soybean seed. Isoflavone O-methyltransferase, catalyzed by isoflavone-O-methyltransferase, is the first step in degradation of daidzein. It may be possible to suppress the activity of this enzyme causing daidzein concentrations to increase yielding higher levels of this beneficial isoflavone in soybean seed.

Some flavonoids have sulfate groups attached to common flavones and flavonols via ester linkages. For example, flavonol 3-sulfotransferase and flavonol 4-sulfotransferase catalyze the stepwise sulfation of quercetin to form quercetin 3,4'-disulfate. The functional significance of flavonoid sulfates in plant tissue is not clear, however, they may play a role

in the detoxification of reactive hydroxyl groups, and/or may be an important molecule for sequestering sulfate ions (Akashi, T. et al. (1998) *Biochemical and Biophysical Research Communications* 251:67-70).

There is a great deal of interest in identifying the genes that encode proteins involved in flavonoid biosynthesis in plants. The genes that code for these enzymes may be used to study the interactions among individuals of the pathways and develop methods to modulate the anthocyanin and flavonol biosynthetic pathways to control flavonoid biosynthesis. Accordingly, the availability of nucleic acid sequences encoding all or a portion of a flavonoid biosynthetic enzyme would facilitate studies to better understand flavonol biosynthesis in plants and provide genetic tools to enhance or otherwise alter flavonol and anthocyanin biosynthesis.

#### SUMMARY OF THE INVENTION

The present invention relates to isolated polynucleotides comprising a nucleotide sequence encoding a polypeptide of at least 348 amino acids that has at least 80% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of soybean isoflavone O-methyltransferase polypeptides of SEQ ID NOs:2, 4, 6, 8 and 10. The present invention also relates to an isolated polynucleotide comprising the complement of the nucleotide sequences described above.

It is preferred that the isolated polynucleotides of the claimed invention consists of a nucleic acid sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7 and 9 that codes for the polypeptide selected from the group consisting of SEQ ID NOs:2, 4, 6, 8 and 10. The present invention also relates to an isolated polynucleotide comprising a nucleotide sequences of at least one of 60 (preferably at least one of 40, most preferably at least one of 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7 and 9 and the complement of such nucleotide sequences.

The present invention relates to a chimeric gene comprising an isolated polynucleotide of the present invention operably linked to suitable regulatory sequences.

The present invention relates to an isolated host cell comprising a chimeric gene of the present invention or an isolated polynucleotide of the present invention. The host cell may be eukaryotic, such as a yeast or a plant cell, or prokaryotic, such as a bacterial cell. The present invention also relates to a virus, preferably a baculovirus, comprising an isolated polynucleotide of the present invention or a chimeric gene of the present invention.

The present invention relates to a process for producing an isolated host cell comprising a chimeric gene of the present invention or an isolated polynucleotide of the present invention, the process comprising either transforming or transfecting an isolated compatible host cell with a chimeric gene or isolated polynucleotide of the present invention.

The present invention relates to an isoflavone O-methyltransferase polypeptide of at least 348 amino acids comprising at least 80% homology based on the Clustal method of alignment compared to a polypeptide selected from the group consisting of SEQ ID NOs:2, 4, 6, 8 and 10.

5       The present invention relates to a method of selecting an isolated polynucleotide that affects the level of expression of an isoflavone O-methyltransferase polypeptide in a host cell, preferably a plant cell, the method comprising the steps of: (a) constructing an isolated polynucleotide of the present invention or an isolated chimeric gene of the present invention; (b) introducing the isolated polynucleotide or the isolated chimeric gene into a host cell;  
10       (c) measuring the level of an isoflavone O-methyltransferase polypeptide in the host cell containing the isolated polynucleotide; and (d) comparing the level of an isoflavone O-methyltransferase polypeptide in the host cell containing the isolated polynucleotide with the level of an isoflavone O-methyltransferase polypeptide in the host cell that does not contain the isolated polynucleotide.

15       The present invention relates to a method of obtaining a nucleic acid fragment encoding a substantial portion of an isoflavone O-methyltransferase polypeptide gene, preferably a plant isoflavone O-methyltransferase polypeptide gene, comprising the steps of: synthesizing an oligonucleotide primer comprising a nucleotide sequence of at least one of 60 (preferably at least one of 40, most preferably at least one of 30) contiguous nucleotides  
20       derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9 and the complement of such nucleotide sequences; and amplifying a nucleic acid fragment (preferably a cDNA inserted in a cloning vector) using the oligonucleotide primer. The amplified nucleic acid fragment preferably will encode a portion of an isoflavone O-methyltransferase amino acid sequence.

25       The present invention also relates to a method of obtaining a nucleic acid fragment encoding all or a substantial portion of the amino acid sequence encoding an isoflavone O-methyltransferase polypeptide comprising the steps of: probing a cDNA or genomic library with an isolated polynucleotide of the present invention; identifying a DNA clone that hybridizes with an isolated polynucleotide of the present invention; isolating the  
30       identified DNA clone; and sequencing the cDNA or genomic fragment that comprises the isolated DNA clone.

The present invention relates to a composition, such as a hybridization mixture, comprising an isolated polynucleotide of the present invention.

35       The present invention relates to an isolated polynucleotide of the present invention comprising at least one of 30 contiguous nucleotides derived from a nucleic acid sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7 and 9.

The present invention relates to an expression cassette comprising an isolated polynucleotide of the present invention operably linked to a promoter.

The present invention relates to a method for positive selection of a transformed cell comprising: (a) transforming a host cell with the chimeric gene of the present invention or an expression cassette of the present invention; and (b) growing the transformed host cell, preferably plant cell, such as a monocot or a dicot, under conditions which allow expression of the isoflavone O-methyltransferase polynucleotide in an amount sufficient to complement a mutant cell and provide altered isoflavone levels to provide a positive selection means.

### BRIEF DESCRIPTION OF THE SEQUENCE DESCRIPTIONS

The invention can be more fully understood from the following detailed description and the accompanying Sequence Listing which form a part of this application.

Table 1 lists the polypeptides that are described herein, the designation of the cDNA clones that comprise the nucleic acid fragments encoding polypeptides representing all or a substantial portion of these polypeptides, and the corresponding identifier (SEQ ID NO:) as used in the attached Sequence Listing. Table 1 also identifies the cDNA clones as individual ESTs ("EST"), the sequences of the entire cDNA inserts comprising the indicated cDNA clones ("FIS"), contigs assembled from two or more ESTs ("Contig"), contigs assembled from an FIS and one or more ESTs ("Contig\*"), or sequences encoding the entire protein derived from an FIS, a contig, or an FIS and PCR ("CGS"). Nucleotide sequence, SEQ ID NO:5 and amino acid sequence SEQ ID NO:6 were determined by further sequence analysis of cDNA clone encoding the amino acid sequence set forth in SEQ ID NO:10. Nucleotide SEQ ID NO:9 and amino acid SEQ ID NO:10 were presented in a U.S. Provisional Application No. 60/113,190, filed December 21, 1998.

The sequence descriptions and Sequence Listing attached hereto comply with the rules governing nucleotide and/or amino acid sequence disclosures in patent applications as set forth in 37 C.F.R. §1.821-1.825.

TABLE 1

#### Flavonoid Biosynthetic Enzymes

Protein	Clone Designation	SEQ ID NO:	
		(Nucleotide)	(Amino Acid)
Isoflavone O-Methyltransferase	se4.pk0007.a11 (FIS)	1	2
Isoflavone O-Methyltransferase	sr1.pk0015.b4 (FIS)	3	4
Isoflavone O-Methyltransferase	srr1c.pk001.b16 (FIS)	5	6
Isoflavone O-Methyltransferase	ssl1c.pk002.d1 (FIS)	7	8
Isoflavone O-Methyltransferase	srr1c.pk001.b16	9	10

The Sequence Listing contains the one letter code for nucleotide sequence characters and the three letter codes for amino acids as defined in conformity with the IUPAC-IUBMB standards described in *Nucleic Acids Res.* 13:3021-3030 (1985) and in the *Biochemical J.*

219 (No. 2):345-373 (1984) which are herein incorporated by reference. The symbols and format used for nucleotide and amino acid sequence data comply with the rules set forth in 37 C.F.R. §1.822.

#### DETAILED DESCRIPTION OF THE INVENTION

5 In the context of this disclosure, a number of terms shall be utilized. As used herein, a "polynucleotide" is a nucleotide sequence such as a nucleic acid fragment. A polynucleotide may be a polymer of RNA or DNA that is single- or double-stranded, that optionally contains synthetic, non-natural or altered nucleotide bases. A polynucleotide in the form of a polymer of DNA may be comprised of one or more segments of cDNA, genomic DNA,  
10 synthetic DNA, or mixtures thereof. An isolated polynucleotide of the present invention may include at least one of 60 contiguous nucleotides, preferably at least one of 40 contiguous nucleotides, most preferably one of at least 30 contiguous nucleotides derived from SEQ ID NOs:1, 3, 5, 7, 9 or the complement of such sequences.

As used herein, "substantially similar" refers to nucleic acid fragments wherein  
15 changes in one or more nucleotide bases results in substitution of one or more amino acids, but do not affect the functional properties of the polypeptide encoded by the nucleotide sequence. "Substantially similar" also refers to nucleic acid fragments wherein changes in one or more nucleotide bases does not affect the ability of the nucleic acid fragment to mediate alteration of gene expression by gene silencing through for example antisense or co-  
20 suppression technology. "Substantially similar" also refers to modifications of the nucleic acid fragments of the instant invention such as deletion or insertion of one or more nucleotides that do not substantially affect the functional properties of the resulting transcript vis-à-vis the ability to mediate gene silencing or alteration of the functional properties of the resulting protein molecule. It is therefore understood that the invention  
25 encompasses more than the specific exemplary nucleotide or amino acid sequences and includes functional equivalents thereof.

Substantially similar nucleic acid fragments may be selected by screening nucleic acid fragments representing subfragments or modifications of the nucleic acid fragments of the instant invention, wherein one or more nucleotides are substituted, deleted and/or inserted,  
30 for their ability to affect the level of the polypeptide encoded by the unmodified nucleic acid fragment in a plant or plant cell. For example, a substantially similar nucleic acid fragment representing at least one of 30 contiguous nucleotides derived from the instant nucleic acid fragment can be constructed and introduced into a plant or plant cell. The level of the polypeptide encoded by the unmodified nucleic acid fragment present in a plant or plant cell  
35 exposed to the substantially similar nucleic fragment can then be compared to the level of the polypeptide in a plant or plant cell that is not exposed to the substantially similar nucleic acid fragment.

For example, it is well known in the art that antisense suppression and co-suppression of gene expression may be accomplished using nucleic acid fragments representing less than the entire coding region of a gene, and by nucleic acid fragments that do not share 100% sequence identity with the gene to be suppressed. Moreover, alterations in a nucleic acid fragment which result in the production of a chemically equivalent amino acid at a given site, but do not effect the functional properties of the encoded polypeptide, are well known in the art. Thus, a codon for the amino acid alanine, a hydrophobic amino acid, may be substituted by a codon encoding another less hydrophobic residue, such as glycine, or a more hydrophobic residue, such as valine, leucine, or isoleucine. Similarly, changes which result in substitution of one negatively charged residue for another, such as aspartic acid for glutamic acid, or one positively charged residue for another, such as lysine for arginine, can also be expected to produce a functionally equivalent product. Nucleotide changes which result in alteration of the N-terminal and C-terminal portions of the polypeptide molecule would also not be expected to alter the activity of the polypeptide. Each of the proposed modifications is well within the routine skill in the art, as is determination of retention of biological activity of the encoded products. Consequently, an isolated polynucleotide comprising a nucleotide sequence of at least one of 60 (preferably at least one of 40, most preferably at least one of 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9 and the complement of such nucleotide sequences may be used in methods of selecting an isolated polynucleotide that affects the expression of a polypeptide (isoflavone O-methyltransferase) in a host cell. A method of selecting an isolated polynucleotide that affects the level of expression of a polypeptide in a host cell (eukaryotic, such as plant or yeast, prokaryotic such as bacterial, or viral) may comprise the steps of: constructing an isolated polynucleotide of the present invention or an isolated chimeric gene of the present invention; introducing the isolated polynucleotide or the isolated chimeric gene into a host cell; measuring the level a polypeptide in the host cell containing the isolated polynucleotide; and comparing the level of a polypeptide in the host cell containing the isolated polynucleotide with the level of a polypeptide in a host cell that does not contain the isolated polynucleotide.

Moreover, substantially similar nucleic acid fragments may also be characterized by their ability to hybridize. Estimates of such homology are provided by either DNA-DNA or DNA-RNA hybridization under conditions of stringency as is well understood by those skilled in the art (Hames and Higgins, Eds. (1985) *Nucleic Acid Hybridisation*, IRL Press, Oxford, U.K.). Stringency conditions can be adjusted to screen for moderately similar fragments, such as homologous sequences from distantly related organisms, to highly similar fragments, such as genes that duplicate functional enzymes from closely related organisms. Post-hybridization washes determine stringency conditions. One set of preferred conditions uses a series of washes starting with 6X SSC, 0.5% SDS at room temperature for 15 min.



then repeated with 2X SSC, 0.5% SDS at 45°C for 30 min, and then repeated twice with 0.2X SSC, 0.5% SDS at 50°C for 30 min. A more preferred set of stringent conditions uses higher temperatures in which the washes are identical to those above except for the temperature of the final two 30 min washes in 0.2X SSC, 0.5% SDS was increased to 60°C.

- 5 Another preferred set of highly stringent conditions uses two final washes in 0.1X SSC, 0.1% SDS at 65°C.

Substantially similar nucleic acid fragments of the instant invention may also be characterized by the percent identity of the amino acid sequences that they encode to the amino acid sequences disclosed herein, as determined by algorithms commonly employed by those skilled in this art. Suitable nucleic acid fragments (isolated polynucleotides of the present invention) encode polypeptides that are at least about 70% identical, preferably at least about 80% identical to the amino acid sequences reported herein. Preferred nucleic acid fragments encode amino acid sequences that are about 85% identical to the amino acid sequences reported herein. More preferred nucleic acid fragments encode amino acid sequences that are at least about 90% identical to the amino acid sequences reported herein. Most preferred are nucleic acid fragments that encode amino acid sequences that are at least about 95% identical to the amino acid sequences reported herein. Suitable nucleic acid fragments not only have the above homologies but typically encode a polypeptide having at least about 50 amino acids, preferably at least about 100 amino acids, more preferably at least about 150 amino acids, still more preferably at least about 200 amino acids, and most preferably at least about 250 amino acids. Sequence alignments and percent identity calculations were performed using the Megalign program of the LASERGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). Multiple alignment of the sequences was performed using the Clustal method of alignment (Higgins and Sharp (1989) *CABIOS*. 5:151-153) with the default parameters (GAP PENALTY=10, GAP LENGTH PENALTY=10). Default parameters for pairwise alignments using the Clustal method were KTUPLE 1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5.

A "substantial portion" of an amino acid or nucleotide sequence comprises an amino acid or a nucleotide sequence that is sufficient to afford putative identification of the protein or gene that the amino acid or nucleotide sequence comprises. Amino acid and nucleotide sequences can be evaluated either manually by one skilled in the art, or by using computer-based sequence comparison and identification tools that employ algorithms such as BLAST (Basic Local Alignment Search Tool; Altschul et al. (1993) *J. Mol. Biol.* 215:403-410; see also [www.ncbi.nlm.nih.gov/BLAST/](http://www.ncbi.nlm.nih.gov/BLAST/)). In general, a sequence of ten or more contiguous amino acids or thirty or more contiguous nucleotides is necessary in order to putatively identify a polypeptide or nucleic acid sequence as homologous to a known protein or gene. Moreover, with respect to nucleotide sequences, gene-specific oligonucleotide probes comprising 30 or more contiguous nucleotides may be used in sequence-dependent methods

of gene identification (e.g., Southern hybridization) and isolation (e.g., *in situ* hybridization of bacterial colonies or bacteriophage plaques). In addition, short oligonucleotides of 12 or more nucleotides may be used as amplification primers in PCR in order to obtain a particular nucleic acid fragment comprising the primers. Accordingly, a "substantial portion" of a nucleotide sequence comprises a nucleotide sequence that will afford specific identification and/or isolation of a nucleic acid fragment comprising the sequence. The instant specification teaches amino acid and nucleotide sequences encoding polypeptides that comprise one or more particular plant proteins. The skilled artisan, having the benefit of the sequences as reported herein, may now use all or a substantial portion of the disclosed sequences for purposes known to those skilled in this art. Accordingly, the instant invention comprises the complete sequences as reported in the accompanying Sequence Listing, as well as substantial portions of those sequences as defined above.

"Codon degeneracy" refers to divergence in the genetic code permitting variation of the nucleotide sequence without effecting the amino acid sequence of an encoded polypeptide. Accordingly, the instant invention relates to any nucleic acid fragment comprising a nucleotide sequence that encodes all or a substantial portion of the amino acid sequences set forth herein. The skilled artisan is well aware of the "codon-bias" exhibited by a specific host cell in usage of nucleotide codons to specify a given amino acid. Therefore, when synthesizing a nucleic acid fragment for improved expression in a host cell, it is desirable to design the nucleic acid fragment such that its frequency of codon usage approaches the frequency of preferred codon usage of the host cell.

"Synthetic nucleic acid fragments" can be assembled from oligonucleotide building blocks that are chemically synthesized using procedures known to those skilled in the art. These building blocks are ligated and annealed to form larger nucleic acid fragments which may then be enzymatically assembled to construct the entire desired nucleic acid fragment. "Chemically synthesized", as related to nucleic acid fragment, means that the component nucleotides were assembled *in vitro*. Manual chemical synthesis of nucleic acid fragments may be accomplished using well established procedures, or automated chemical synthesis can be performed using one of a number of commercially available machines. Accordingly, the nucleic acid fragments can be tailored for optimal gene expression based on optimization of nucleotide sequence to reflect the codon bias of the host cell. The skilled artisan appreciates the likelihood of successful gene expression if codon usage is biased towards those codons favored by the host. Determination of preferred codons can be based on a survey of genes derived from the host cell where sequence information is available.

"Gene" refers to a nucleic acid fragment that expresses a specific protein, including regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding sequences) the coding sequence. "Native gene" refers to a gene as found in nature with its own regulatory sequences. "Chimeric gene" refers any gene that is not a native gene,

comprising regulatory and coding sequences that are not found together in nature.

Accordingly, a chimeric gene may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that found in nature.

5 "Endogenous gene" refers to a native gene in its natural location in the genome of an organism. A "foreign" gene refers to a gene not normally found in the host organism, but that is introduced into the host organism by gene transfer. Foreign genes can comprise native genes inserted into a non-native organism, or chimeric genes. A "transgene" is a gene that has been introduced into the genome by a transformation procedure.

10 "Coding sequence" refers to a nucleotide sequence that codes for a specific amino acid sequence. "Regulatory sequences" refer to nucleotide sequences located upstream (5' non-coding sequences), within, or downstream (3' non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the associated coding sequence. Regulatory sequences may include promoters, translation  
15 leader sequences, introns, and polyadenylation recognition sequences.

"Promoter" refers to a nucleotide sequence capable of controlling the expression of a coding sequence or functional RNA. In general, a coding sequence is located 3' to a promoter sequence. The promoter sequence consists of proximal and more distal upstream elements, the latter elements often referred to as enhancers. Accordingly, an "enhancer" is a  
20 nucleotide sequence which can stimulate promoter activity and may be an innate element of the promoter or a heterologous element inserted to enhance the level or tissue-specificity of a promoter. Promoters may be derived in their entirety from a native gene, or be composed of different elements derived from different promoters found in nature, or even comprise synthetic nucleotide segments. It is understood by those skilled in the art that different  
25 promoters may direct the expression of a gene in different tissues or cell types, or at different stages of development, or in response to different environmental conditions. Promoters which cause a nucleic acid fragment to be expressed in most cell types at most times are commonly referred to as "constitutive promoters". New promoters of various types useful in plant cells are constantly being discovered; numerous examples may be  
30 found in the compilation by Okamuro and Goldberg (1989) *Biochemistry of Plants* 15:1-82. It is further recognized that since in most cases the exact boundaries of regulatory sequences have not been completely defined, nucleic acid fragments of different lengths may have identical promoter activity.

The "translation leader sequence" refers to a nucleotide sequence located between the  
35 promoter sequence of a gene and the coding sequence. The translation leader sequence is present in the fully processed mRNA upstream of the translation start sequence. The translation leader sequence may affect processing of the primary transcript to mRNA.

mRNA stability or translation efficiency. Examples of translation leader sequences have been described (Turner and Foster (1995) *Mol. Biotechnol.* 3:225-236).

The "3' non-coding sequences" refer to nucleotide sequences located downstream of a coding sequence and include polyadenylation recognition sequences and other sequences encoding regulatory signals capable of affecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by affecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor. The use of different 3' non-coding sequences is exemplified by Ingelbrecht et al. (1989) *Plant Cell* 1:671-680.

"RNA transcript" refers to the product resulting from RNA polymerase-catalyzed transcription of a DNA sequence. When the RNA transcript is a perfect complementary copy of the DNA sequence, it is referred to as the primary transcript or it may be a RNA sequence derived from posttranscriptional processing of the primary transcript and is referred to as the mature RNA. "Messenger RNA (mRNA)" refers to the RNA that is without introns and that can be translated into polypeptide by the cell. "cDNA" refers to a double-stranded DNA that is complementary to and derived from mRNA. "Sense" RNA refers to an RNA transcript that includes the mRNA and so can be translated into a polypeptide by the cell. "Antisense RNA" refers to an RNA transcript that is complementary to all or part of a target primary transcript or mRNA and that blocks the expression of a target gene (see U.S. Patent No. 5,107,065, incorporated herein by reference). The complementarity of an antisense RNA may be with any part of the specific nucleotide sequence, i.e., at the 5' non-coding sequence, 3' non-coding sequence, introns, or the coding sequence. "Functional RNA" refers to sense RNA, antisense RNA, ribozyme RNA, or other RNA that may not be translated but yet has an effect on cellular processes.

The term "operably linked" refers to the association of two or more nucleic acid fragments on a single nucleic acid fragment so that the function of one is affected by the other. For example, a promoter is operably linked with a coding sequence when it is capable of affecting the expression of that coding sequence (i.e., that the coding sequence is under the transcriptional control of the promoter). Coding sequences can be operably linked to regulatory sequences in sense or antisense orientation.

The term "expression", as used herein, refers to the transcription and stable accumulation of sense (mRNA) or antisense RNA derived from the nucleic acid fragment of the invention. Expression may also refer to translation of mRNA into a polypeptide. "Antisense inhibition" refers to the production of antisense RNA transcripts capable of suppressing the expression of the target protein. "Overexpression" refers to the production of a gene product in transgenic organisms that exceeds levels of production in normal or non-transformed organisms. "Co-suppression" refers to the production of sense RNA transcripts capable of suppressing the expression of identical or substantially similar foreign or endogenous genes (U.S. Patent No. 5,231,020, incorporated herein by reference).

"Altered levels" refers to the production of gene product(s) in transgenic organisms in amounts or proportions that differ from that of normal or non-transformed organisms.

"Mature" protein refers to a post-translationally processed polypeptide; i.e., one from which any pre- or propeptides present in the primary translation product have been removed.

5 "Precursor" protein refers to the primary product of translation of mRNA; i.e., with pre- and propeptides still present. Pre- and propeptides may be but are not limited to intracellular localization signals.

A "chloroplast transit peptide" is an amino acid sequence which is translated in conjunction with a protein and directs the protein to the chloroplast or other plastid types  
10 present in the cell in which the protein is made. "Chloroplast transit sequence" refers to a nucleotide sequence that encodes a chloroplast transit peptide. A "signal peptide" is an amino acid sequence which is translated in conjunction with a protein and directs the protein to the secretory system (Chrispeels (1991) *Ann. Rev. Plant Phys. Plant Mol. Biol.* 42:21-53). If the protein is to be directed to a vacuole, a vacuolar targeting signal (*supra*) can further be  
15 added, or if to the endoplasmic reticulum, an endoplasmic reticulum retention signal (*supra*) may be added. If the protein is to be directed to the nucleus, any signal peptide present should be removed and instead a nuclear localization signal included (Raikhel (1992) *Plant Phys.* 100:1627-1632).

"Transformation" refers to the transfer of a nucleic acid fragment into the genome of a  
20 host organism, resulting in genetically stable inheritance. Host organisms containing the transformed nucleic acid fragments are referred to as "transgenic" organisms. Examples of methods of plant transformation include *Agrobacterium*-mediated transformation (De Blaere et al. (1987) *Meth. Enzymol.* 143:277) and particle-accelerated or "gene gun" transformation technology (Klein et al. (1987) *Nature (London)* 327:70-73; U.S. Patent No. 4,945,050,  
25 incorporated herein by reference).

Standard recombinant DNA and molecular cloning techniques used herein are well known in the art and are described more fully in Sambrook et al. *Molecular Cloning: A Laboratory Manual*; Cold Spring Harbor Laboratory Press: Cold Spring Harbor, 1989 (hereinafter "Maniatis").

30 Nucleic acid fragments encoding at least a portion of several flavonoid biosynthetic enzymes have been isolated and identified by comparison of random plant cDNA sequences to public databases containing nucleotide and protein sequences using the BLAST algorithms well known to those skilled in the art. The nucleic acid fragments of the instant invention may be used to isolate cDNAs and genes encoding homologous proteins from the  
35 same or other plant species. Isolation of homologous genes using sequence-dependent protocols is well known in the art. Examples of sequence-dependent protocols include, but are not limited to, methods of nucleic acid hybridization, and methods of DNA and RNA

amplification as exemplified by various uses of nucleic acid amplification technologies (e.g., polymerase chain reaction, ligase chain reaction).

For example, genes encoding other isoflavone O-methyltransferase proteins, either as cDNAs or genomic DNAs, could be isolated directly by using all or a portion of the instant nucleic acid fragments as DNA hybridization probes to screen libraries from any desired plant employing methodology well known to those skilled in the art. Specific oligonucleotide probes based upon the instant nucleic acid sequences can be designed and synthesized by methods known in the art (Maniatis). Moreover, the entire sequences can be used directly to synthesize DNA probes by methods known to the skilled artisan such as random primer DNA labeling, nick translation, or end-labeling techniques, or RNA probes using available *in vitro* transcription systems. In addition, specific primers can be designed and used to amplify a part or all of the instant sequences. The resulting amplification products can be labeled directly during amplification reactions or labeled after amplification reactions, and used as probes to isolate full length cDNA or genomic fragments under conditions of appropriate stringency.

In addition, two short segments of the instant nucleic acid fragments may be used in polymerase chain reaction protocols to amplify longer nucleic acid fragments encoding homologous genes from DNA or RNA. The polymerase chain reaction may also be performed on a library of cloned nucleic acid fragments wherein the sequence of one primer is derived from the instant nucleic acid fragments, and the sequence of the other primer takes advantage of the presence of the polyadenylic acid tracts to the 3' end of the mRNA precursor encoding plant genes. Alternatively, the second primer sequence may be based upon sequences derived from the cloning vector. For example, the skilled artisan can follow the RACE protocol (Frohman et al. (1988) *Proc. Natl. Acad. Sci. USA* 85:8998-9002) to generate cDNAs by using PCR to amplify copies of the region between a single point in the transcript and the 3' or 5' end. Primers oriented in the 3' and 5' directions can be designed from the instant sequences. Using commercially available 3' RACE or 5' RACE systems (BRL), specific 3' or 5' cDNA fragments can be isolated (Ohara et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:5673-5677; Loh et al. (1989) *Science* 243:217-220). Products generated by the 3' and 5' RACE procedures can be combined to generate full-length cDNAs (Frohman and Martin (1989) *Techniques* 1:165). Consequently, a polynucleotide comprising a nucleotide sequence of at least one of 60 (preferably one of at least 40, most preferably one of at least 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9 and the complement of such nucleotide sequences may be used in such methods to obtain a nucleic acid fragment encoding a substantial portion of an amino acid sequence of a polypeptide. The present invention relates to a method of obtaining a nucleic acid fragment encoding a substantial portion of a polypeptide of a gene (such as isoflavone O-methyltransferase) preferably a substantial

portion of a plant polypeptide of a gene, comprising the steps of: synthesizing an oligonucleotide primer comprising a nucleotide sequence of at least one of 60 (preferably at least one of 40, most preferably at least one of 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9 and the complement of such nucleotide sequences; and amplifying a nucleic acid fragment (preferably a cDNA inserted in a cloning vector) using the oligonucleotide primer. The amplified nucleic acid fragment preferably will encode a portion of a polypeptide (such as isoflavone O-methyltransferase).

Availability of the instant nucleotide and deduced amino acid sequences facilitates immunological screening of cDNA expression libraries. Synthetic peptides representing portions of the instant amino acid sequences may be synthesized. These peptides can be used to immunize animals to produce polyclonal or monoclonal antibodies with specificity for peptides or proteins comprising the amino acid sequences. These antibodies can be then be used to screen cDNA expression libraries to isolate full-length cDNA clones of interest (Lerner (1984) *Adv. Immunol.* 36:1-34; Maniatis).

The nucleic acid fragments of the instant invention may be used to create transgenic plants in which the disclosed polypeptides are present at higher or lower levels than normal or in cell types or developmental stages in which they are not normally found. This would have the effect of altering the level isoflavones in those cells.

Overexpression of the proteins of the instant invention may be accomplished by first constructing a chimeric gene in which the coding region is operably linked to a promoter capable of directing expression of a gene in the desired tissues at the desired stage of development. The chimeric gene may comprise promoter sequences and translation leader sequences derived from the same genes. 3' Non-coding sequences encoding transcription termination signals may also be provided. The instant chimeric gene may also comprise one or more introns in order to facilitate gene expression.

Plasmid vectors comprising the isolated polynucleotide (or chimeric gene) may be constructed. The choice of plasmid vector is dependent upon the method that will be used to transform host plants. The skilled artisan is well aware of the genetic elements that must be present on the plasmid vector in order to successfully transform, select and propagate host cells containing the chimeric gene. The skilled artisan will also recognize that different independent transformation events will result in different levels and patterns of expression (Jones et al. (1985) *EMBO J.* 4:2411-2418; De Almeida et al. (1989) *Mol. Gen. Genetics* 218:78-86), and thus that multiple events must be screened in order to obtain lines displaying the desired expression level and pattern. Such screening may be accomplished by Southern analysis of DNA, Northern analysis of mRNA expression, Western analysis of protein expression, or phenotypic analysis.

For some applications it may be useful to direct the instant polypeptides to different cellular compartments, or to facilitate its secretion from the cell. It is thus envisioned that the chimeric gene described above may be further supplemented by directing the coding sequence to encode the instant polypeptides with appropriate intracellular targeting sequences such as transit sequences (Keegstra (1989) *Cell* 56:247-253), signal sequences or sequences encoding endoplasmic reticulum localization (Chrispeels (1991) *Ann. Rev. Plant Phys. Plant Mol. Biol.* 42:21-53), or nuclear localization signals (Raikhel (1992) *Plant Phys.* 100:1627-1632) with or without removing targeting sequences that are already present. While the references cited give examples of each of these, the list is not exhaustive and more targeting signals of use may be discovered in the future.

It may also be desirable to reduce or eliminate expression of genes encoding the instant polypeptides in plants for some applications. In order to accomplish this, a chimeric gene designed for co-suppression of the instant polypeptide can be constructed by linking a gene or gene fragment encoding that polypeptide to plant promoter sequences.

Alternatively, a chimeric gene designed to express antisense RNA for all or part of the instant nucleic acid fragment can be constructed by linking the gene or gene fragment in reverse orientation to plant promoter sequences. Either the co-suppression or antisense chimeric genes could be introduced into plants via transformation wherein expression of the corresponding endogenous genes are reduced or eliminated.

Molecular genetic solutions to the generation of plants with altered gene expression have a decided advantage over more traditional plant breeding approaches. Changes in plant phenotypes can be produced by specifically inhibiting expression of one or more genes by antisense inhibition or cosuppression (U.S. Patent Nos. 5,190,931, 5,107,065 and 5,283,323). An antisense or cosuppression construct would act as a dominant negative regulator of gene activity. While conventional mutations can yield negative regulation of gene activity these effects are most likely recessive. The dominant negative regulation available with a transgenic approach may be advantageous from a breeding perspective. In addition, the ability to restrict the expression of specific phenotype to the reproductive tissues of the plant by the use of tissue specific promoters may confer agronomic advantages relative to conventional mutations which may have an effect in all tissues in which a mutant gene is ordinarily expressed.

The person skilled in the art will know that special considerations are associated with the use of antisense or cosuppression technologies in order to reduce expression of particular genes. For example, the proper level of expression of sense or antisense genes may require the use of different chimeric genes utilizing different regulatory elements known to the skilled artisan. Once transgenic plants are obtained by one of the methods described above, it will be necessary to screen individual transgenics for those that most effectively display the desired phenotype. Accordingly, the skilled artisan will develop methods for screening



large numbers of transformants. The nature of these screens will generally be chosen on practical grounds. For example, one can screen by looking for changes in gene expression by using antibodies specific for the protein encoded by the gene being suppressed, or one could establish assays that specifically measure enzyme activity. A preferred method will be one which allows large numbers of samples to be processed rapidly, since it will be expected that a large number of transformants will be negative for the desired phenotype.

The instant polypeptides (or portions thereof) may be produced in heterologous host cells, particularly in the cells of microbial hosts, and can be used to prepare antibodies to these proteins by methods well known to those skilled in the art. The antibodies are useful for detecting the polypeptides of the instant invention *in situ* in cells or *in vitro* in cell extracts. Preferred heterologous host cells for production of the instant polypeptides are microbial hosts. Microbial expression systems and expression vectors containing regulatory sequences that direct high level expression of foreign proteins are well known to those skilled in the art. Any of these could be used to construct a chimeric gene for production of the instant polypeptides. This chimeric gene could then be introduced into appropriate microorganisms via transformation to provide high level expression of the encoded flavonoid biosynthetic enzyme. An example of a vector for high level expression of the instant polypeptides in a bacterial host is provided (Example 6).

All or a substantial portion of the nucleic acid fragments of the instant invention may also be used as probes for genetically and physically mapping the genes that they are a part of, and as markers for traits linked to those genes. Such information may be useful in plant breeding in order to develop lines with desired phenotypes. For example, the instant nucleic acid fragments may be used as restriction fragment length polymorphism (RFLP) markers. Southern blots (Maniatis) of restriction-digested plant genomic DNA may be probed with the nucleic acid fragments of the instant invention. The resulting banding patterns may then be subjected to genetic analyses using computer programs such as MapMaker (Lander et al. (1987) *Genomics* 1:174-181) in order to construct a genetic map. In addition, the nucleic acid fragments of the instant invention may be used to probe Southern blots containing restriction endonuclease-treated genomic DNAs of a set of individuals representing parent and progeny of a defined genetic cross. Segregation of the DNA polymorphisms is noted and used to calculate the position of the instant nucleic acid sequence in the genetic map previously obtained using this population (Botstein et al. (1980) *Am. J. Hum. Genet.* 32:314-331).

The production and use of plant gene-derived probes for use in genetic mapping is described in Bernatzky and Tanksley (1986) *Plant Mol. Biol. Reporter* 4:37-41. Numerous publications describe genetic mapping of specific cDNA clones using the methodology outlined above or variations thereof. For example, F2 intercross populations, backcross

populations, randomly mated populations, near isogenic lines, and other sets of individuals may be used for mapping. Such methodologies are well known to those skilled in the art.

Nucleic acid probes derived from the instant nucleic acid sequences may also be used for physical mapping (i.e., placement of sequences on physical maps; *see* Hoheisel et al. In: 5 *Nonmammalian Genomic Analysis: A Practical Guide*, Academic press 1996, pp. 319-346, and references cited therein).

In another embodiment, nucleic acid probes derived from the instant nucleic acid sequences may be used in direct fluorescence *in situ* hybridization (FISH) mapping (Trask 10 (1991) *Trends Genet.* 7:149-154). Although current methods of FISH mapping favor use of large clones (several to several hundred KB; *see* Laan et al. (1995) *Genome Res.* 5:13-20), improvements in sensitivity may allow performance of FISH mapping using shorter probes.

A variety of nucleic acid amplification-based methods of genetic and physical mapping may be carried out using the instant nucleic acid sequences. Examples include allele-specific amplification (Kazazian (1989) *J. Lab. Clin. Med.* 11:95-96), polymorphism 15 of PCR-amplified fragments (CAPS; Sheffield et al. (1993) *Genomics* 16:325-332), allele-specific ligation (Landegren et al. (1988) *Science* 241:1077-1080), nucleotide extension reactions (Sokolov (1990) *Nucleic Acid Res.* 18:3671), Radiation Hybrid Mapping (Walter et al. (1997) *Nat. Genet.* 7:22-28) and Happy Mapping (Dear and Cook (1989) *Nucleic Acid Res.* 17:6795-6807). For these methods, the sequence of a nucleic acid fragment is used to 20 design and produce primer pairs for use in the amplification reaction or in primer extension reactions. The design of such primers is well known to those skilled in the art. In methods employing PCR-based genetic mapping, it may be necessary to identify DNA sequence differences between the parents of the mapping cross in the region corresponding to the instant nucleic acid sequence. This, however, is generally not necessary for mapping 25 methods.

Loss of function mutant phenotypes may be identified for the instant cDNA clones either by targeted gene disruption protocols or by identifying specific mutants for these genes contained in a maize population carrying mutations in all possible genes (Ballinger and Benzer (1989) *Proc. Natl. Acad. Sci USA* 86:9402-9406; Koes et al. (1995) *Proc. Natl. Acad. Sci USA* 92:8149-8153; Bensen et al. (1995) *Plant Cell* 7:75-84). The latter approach 30 may be accomplished in two ways. First, short segments of the instant nucleic acid fragments may be used in polymerase chain reaction protocols in conjunction with a mutation tag sequence primer on DNAs prepared from a population of plants in which Mutator transposons or some other mutation-causing DNA element has been introduced (see Bensen, *supra*). The amplification of a specific DNA fragment with these primers 35 indicates the insertion of the mutation tag element in or near the plant gene encoding the instant polypeptides. Alternatively, the instant nucleic acid fragment may be used as a hybridization probe against PCR amplification products generated from the mutation

population using the mutation tag sequence primer in conjunction with an arbitrary genomic site primer, such as that for a restriction enzyme site-anchored synthetic adaptor. With either method, a plant containing a mutation in the endogenous gene encoding the instant polypeptides can be identified and obtained. This mutant plant can then be used to determine or confirm the natural function of the instant polypeptides disclosed herein.

### EXAMPLES

The present invention is further defined in the following Examples, in which all parts and percentages are by weight and degrees are Celsius, unless otherwise stated. It should be understood that these Examples, while indicating preferred embodiments of the invention, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.

#### EXAMPLE 1

##### Composition of cDNA Libraries; Isolation and Sequencing of cDNA Clones

cDNA libraries representing mRNAs from various soybean tissues were prepared. The characteristics of the libraries are described below.

TABLE 2

cDNA Libraries from Soybean

Library	Tissue	Clone
se4	Soybean Embryo, 19 Days After Flowering	se4.pk0007.a11
sr1	Soybean Root	sr1.pk0015.b4
srr1c	Soybean 8-Day-Old Root	srr1c.pk001.b16
ssl1c	Soybean Seed 25 Days After Flowering	ssl1c.pk002.d1

cDNA libraries may be prepared by any one of many methods available. For example, the cDNAs may be introduced into plasmid vectors by first preparing the cDNA libraries in Uni-ZAP™ XR vectors according to the manufacturer's protocol (Stratagene Cloning Systems, La Jolla, CA). The Uni-ZAP™ XR libraries are converted into plasmid libraries according to the protocol provided by Stratagene. Upon conversion, cDNA inserts will be contained in the plasmid vector pBluescript. In addition, the cDNAs may be introduced directly into precut Bluescript II SK(+) vectors (Stratagene) using T4 DNA ligase (New England Biolabs), followed by transfection into DH10B cells according to the manufacturer's protocol (GIBCO BRL Products). Once the cDNA inserts are in plasmid vectors, plasmid DNAs are prepared from randomly picked bacterial colonies containing recombinant pBluescript plasmids, or the insert cDNA sequences are amplified via polymerase chain reaction using primers specific for vector sequences flanking the inserted

cDNA sequences. Amplified insert DNAs or plasmid DNAs are sequenced in dye-primer sequencing reactions to generate partial cDNA sequences (expressed sequence tags or "ESTs"; see Adams et al., (1991) *Science* 252:1651-1656). The resulting ESTs are analyzed using a Perkin Elmer Model 377 fluorescent sequencer.

5

## EXAMPLE 2

### Identification of cDNA Clones

cDNA clones encoding flavonoid biosynthetic enzymes were identified by conducting BLAST (Basic Local Alignment Search Tool; Altschul et al. (1993) *J. Mol. Biol.* 215:403-410; see also [www.ncbi.nlm.nih.gov/BLAST/](http://www.ncbi.nlm.nih.gov/BLAST/)) searches for similarity to sequences contained in the BLAST "nr" database (comprising all non-redundant GenBank CDS translations, sequences derived from the 3-dimensional structure Brookhaven Protein Data Bank, the last major release of the SWISS-PROT protein sequence database, EMBL, and DDBJ databases). The cDNA sequences obtained in Example 1 were analyzed for similarity to all publicly available DNA sequences contained in the "nr" database using the BLASTN algorithm provided by the National Center for Biotechnology Information (NCBI). The DNA sequences were translated in all reading frames and compared for similarity to all publicly available protein sequences contained in the "nr" database using the BLASTX algorithm (Gish and States (1993) *Nat. Genet.* 3:266-272) provided by the NCBI. For convenience, the P-value (probability) of observing a match of a cDNA sequence to a sequence contained in the searched databases merely by chance as calculated by BLAST are reported herein as "pLog" values, which represent the negative of the logarithm of the reported P-value. Accordingly, the greater the pLog value, the greater the likelihood that the cDNA sequence and the BLAST "hit" represent homologous proteins.

10

15

20

25

## EXAMPLE 3

### Characterization of cDNA Clones Encoding Isoflavone O-Methyltransferase

The BLASTX search using the EST sequences from clones listed in Table 3 revealed similarity of the polypeptides encoded by the cDNAs to isoflavone O-methyltransferase from *Medicago sativa* (NCBI Identifier No. gi 2580582) and *Medicago sativa* (NCBI Identifier No. gi 2580584). Shown in Table 3 are the BLAST results for individual ESTs ("EST"), the sequences of the entire cDNA inserts comprising the indicated cDNA clones ("FIS"), contigs assembled from two or more ESTs ("Contig"), contigs assembled from an FIS and one or more ESTs ("Contig\*"), or sequences encoding the entire protein derived from an FIS, a contig, or an FIS and PCR ("CGS"):

30

TABLE 3

BLAST Results for Sequences Encoding Polypeptides Homologous  
to *Medicago sativa* Isoflavone O-Methyltransferase

Clone	Status	BLAST pLog Score
se4.pk0007.a11	(FIS)	117.00 (gi 2580582)
sr1.pk0015.b4	(FIS)	142.00 (gi 2580584)
srr1c.pk001.b16	(FIS)	122.00 (gi 2580582)
ssl1c.pk002.d1	(FIS)	118.00 (gi 2580582)

- 5 The data in Table 4 represents a calculation of the percent identity of the amino acid sequences set forth in SEQ ID NOs:2, 4, 6 and 8 and the *Medicago sativa* sequences.

TABLE 4

10 Percent Identity of Amino Acid Sequences Deduced From the Nucleotide Sequences of cDNA Clones Encoding Polypeptides Homologous to *Medicago sativa* Isoflavone O-Methyltransferase

SEQ ID NO.	Percent Identity to
2	55% (gi 2580582)
4	66% (gi 2580584)
6	58% (gi 2580582)
8	56% (gi 2580582)

- Sequence alignments and percent identity calculations were performed using the Megalign program of the LASERGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). Multiple alignment of the sequences was performed using the Clustal method of alignment (Higgins and Sharp (1989) *CABIOS*. 5:151-153) with the default parameters (GAP PENALTY=10, GAP LENGTH PENALTY=10). Default parameters for pairwise alignments using the Clustal method were KTUPLE 1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5. Sequence alignments and BLAST scores and probabilities indicate that the nucleic acid fragments comprising the instant cDNA clones encode a substantial portion of a isoflavone O-methyltransferase. These sequences represent the first soybean sequences encoding isoflavone O-methyltransferase.

EXAMPLE 4Expression of Chimeric Genes in Monocot Cells

- 25 A chimeric gene comprising a cDNA encoding the instant polypeptides in sense orientation with respect to the maize 27 kD zein promoter that is located 5' to the cDNA fragment, and the 10 kD zein 3' end that is located 3' to the cDNA fragment, can be constructed. The cDNA fragment of this gene may be generated by polymerase chain reaction (PCR) of the cDNA clone using appropriate oligonucleotide primers. Cloning sites

(NcoI or SmaI) can be incorporated into the oligonucleotides to provide proper orientation of the DNA fragment when inserted into the digested vector pML103 as described below. Amplification is then performed in a standard PCR. The amplified DNA is then digested with restriction enzymes NcoI and SmaI and fractionated on an agarose gel. The appropriate  
5 band can be isolated from the gel and combined with a 4.9 kb NcoI-SmaI fragment of the plasmid pML103. Plasmid pML103 has been deposited under the terms of the Budapest Treaty at ATCC (American Type Culture Collection, 10801 University Blvd., Manassas, VA 20110-2209), and bears accession number ATCC 97366. The DNA segment from pML103 contains a 1.05 kb SalI-NcoI promoter fragment of the maize 27 kD zein gene and  
10 a 0.96 kb SmaI-SalI fragment from the 3' end of the maize 10 kD zein gene in the vector pGem9Zf(+) (Promega). Vector and insert DNA can be ligated at 15°C overnight, essentially as described (Maniatis). The ligated DNA may then be used to transform *E. coli* XL1-Blue (Epicurian Coli XL-1 Blue™; Stratagene). Bacterial transformants can be screened by restriction enzyme digestion of plasmid DNA and limited nucleotide sequence  
15 analysis using the dideoxy chain termination method (Sequenase™ DNA Sequencing Kit; U.S. Biochemical). The resulting plasmid construct would comprise a chimeric gene encoding, in the 5' to 3' direction, the maize 27 kD zein promoter, a cDNA fragment encoding the instant polypeptides, and the 10 kD zein 3' region.

The chimeric gene described above can then be introduced into corn cells by the  
20 following procedure. Immature corn embryos can be dissected from developing caryopses derived from crosses of the inbred corn lines H99 and LH132. The embryos are isolated 10 to 11 days after pollination when they are 1.0 to 1.5 mm long. The embryos are then placed with the axis-side facing down and in contact with agarose-solidified N6 medium (Chu et al. (1975) *Sci. Sin. Peking* 18:659-668). The embryos are kept in the dark at 27°C. Friable  
25 embryogenic callus consisting of undifferentiated masses of cells with somatic proembryoids and embryoids borne on suspensor structures proliferates from the scutellum of these immature embryos. The embryogenic callus isolated from the primary explant can be cultured on N6 medium and sub-cultured on this medium every 2 to 3 weeks.

The plasmid, p35S/Ac (obtained from Dr. Peter Eckes, Hoechst Ag, Frankfurt, Germany) may be used in transformation experiments in order to provide for a selectable  
30 marker. This plasmid contains the *Pat* gene (see European Patent Publication 0 242 236) which encodes phosphinothricin acetyl transferase (PAT). The enzyme PAT confers resistance to herbicidal glutamine synthetase inhibitors such as phosphinothricin. The *pat* gene in p35S/Ac is under the control of the 35S promoter from Cauliflower Mosaic Virus  
35 (Odell et al. (1985) *Nature* 313:810-812) and the 3' region of the nopaline synthase gene from the T-DNA of the Ti plasmid of *Agrobacterium tumefaciens*.

The particle bombardment method (Klein et al. (1987) *Nature* 327:70-73) may be used to transfer genes to the callus culture cells. According to this method, gold particles (1 µm

in diameter) are coated with DNA using the following technique. Ten  $\mu\text{g}$  of plasmid DNAs are added to 50  $\mu\text{L}$  of a suspension of gold particles (60 mg per mL). Calcium chloride (50  $\mu\text{L}$  of a 2.5 M solution) and spermidine free base (20  $\mu\text{L}$  of a 1.0 M solution) are added to the particles. The suspension is vortexed during the addition of these solutions. After  
5 10 minutes, the tubes are briefly centrifuged (5 sec at 15,000 rpm) and the supernatant removed. The particles are resuspended in 200  $\mu\text{L}$  of absolute ethanol, centrifuged again and the supernatant removed. The ethanol rinse is performed again and the particles resuspended in a final volume of 30  $\mu\text{L}$  of ethanol. An aliquot (5  $\mu\text{L}$ ) of the DNA-coated  
10 gold particles can be placed in the center of a Kapton<sup>TM</sup> flying disc (Bio-Rad Labs). The particles are then accelerated into the corn tissue with a Biolistic<sup>TM</sup> PDS-1000/He (Bio-Rad Instruments, Hercules CA), using a helium pressure of 1000 psi, a gap distance of 0.5 cm and a flying distance of 1.0 cm.

For bombardment, the embryogenic tissue is placed on filter paper over agarose-solidified N6 medium. The tissue is arranged as a thin lawn and covered a circular area of  
15 about 5 cm in diameter. The petri dish containing the tissue can be placed in the chamber of the PDS-1000/He approximately 8 cm from the stopping screen. The air in the chamber is then evacuated to a vacuum of 28 inches of Hg. The macrocarrier is accelerated with a helium shock wave using a rupture membrane that bursts when the He pressure in the shock tube reaches 1000 psi.

20 Seven days after bombardment the tissue can be transferred to N6 medium that contains gluphosinate (2 mg per liter) and lacks casein or proline. The tissue continues to grow slowly on this medium. After an additional 2 weeks the tissue can be transferred to fresh N6 medium containing gluphosinate. After 6 weeks, areas of about 1 cm in diameter of actively growing callus can be identified on some of the plates containing the glufosinate-  
25 supplemented medium. These calli may continue to grow when sub-cultured on the selective medium.

Plants can be regenerated from the transgenic callus by first transferring clusters of tissue to N6 medium supplemented with 0.2 mg per liter of 2,4-D. After two weeks the tissue can be transferred to regeneration medium (Fromm et al. (1990) *Bio/Technology*  
30 8:833-839).

### EXAMPLE 5

#### Expression of Chimeric Genes in Dicot Cells

A seed-specific expression cassette composed of the promoter and transcription terminator from the gene encoding the  $\beta$  subunit of the seed storage protein phaseolin from  
35 the bean *Phaseolus vulgaris* (Doyle et al. (1986) *J. Biol. Chem.* 261:9228-9238) can be used for expression of the instant polypeptides in transformed soybean. The phaseolin cassette includes about 500 nucleotides upstream (5') from the translation initiation codon and about 1650 nucleotides downstream (3') from the translation stop codon of phaseolin. Between the

5' and 3' regions are the unique restriction endonuclease sites Nco I (which includes the ATG translation initiation codon), Sma I, Kpn I and Xba I. The entire cassette is flanked by Hind III sites.

The cDNA fragment of this gene may be generated by polymerase chain reaction (PCR) of the cDNA clone using appropriate oligonucleotide primers. Cloning sites can be incorporated into the oligonucleotides to provide proper orientation of the DNA fragment when inserted into the expression vector. Amplification is then performed as described above, and the isolated fragment is inserted into a pUC18 vector carrying the seed expression cassette.

Soybean embryos may then be transformed with the expression vector comprising sequences encoding the instant polypeptides. To induce somatic embryos, cotyledons, 3-5 mm in length dissected from surface sterilized, immature seeds of the soybean cultivar A2872, can be cultured in the light or dark at 26°C on an appropriate agar medium for 6-10 weeks. Somatic embryos which produce secondary embryos are then excised and placed into a suitable liquid medium. After repeated selection for clusters of somatic embryos which multiplied as early, globular staged embryos, the suspensions are maintained as described below.

Soybean embryogenic suspension cultures can maintained in 35 mL liquid media on a rotary shaker, 150 rpm, at 26°C with florescent lights on a 16:8 hour day/night schedule.

Cultures are subcultured every two weeks by inoculating approximately 35 mg of tissue into 35 mL of liquid medium.

Soybean embryogenic suspension cultures may then be transformed by the method of particle gun bombardment (Klein et al. (1987) *Nature* (London) 327:70-73, U.S. Patent No. 4,945,050). A DuPont Biolistic™ PDS1000/HE instrument (helium retrofit) can be used for these transformations.

A selectable marker gene which can be used to facilitate soybean transformation is a chimeric gene composed of the 35S promoter from Cauliflower Mosaic Virus (Odell et al. (1985) *Nature* 313:810-812), the hygromycin phosphotransferase gene from plasmid pJR225 (from *E. coli*; Gritz et al.(1983) *Gene* 25:179-188) and the 3' region of the nopaline synthase gene from the T-DNA of the Ti plasmid of *Agrobacterium tumefaciens*. The seed expression cassette comprising the phaseolin 5' region, the fragment encoding the instant polypeptides and the phaseolin 3' region can be isolated as a restriction fragment. This fragment can then be inserted into a unique restriction site of the vector carrying the marker gene.

To 50 µL of a 60 mg/mL 1 µm gold particle suspension is added (in order): 5 µL DNA (1 µg/µL), 20 µl spermidine (0.1 M), and 50 µL CaCl<sub>2</sub> (2.5 M). The particle preparation is then agitated for three minutes, spun in a microfuge for 10 seconds and the supernatant removed. The DNA-coated particles are then washed once in 400 µL 70% ethanol and resuspended in 40 µL of anhydrous ethanol. The DNA/particle suspension can



be sonicated three times for one second each. Five  $\mu$ L of the DNA-coated gold particles are then loaded on each macro carrier disk.

Approximately 300-400 mg of a two-week-old suspension culture is placed in an empty 60x15 mm petri dish and the residual liquid removed from the tissue with a pipette.

5 For each transformation experiment, approximately 5-10 plates of tissue are normally bombarded. Membrane rupture pressure is set at 1100 psi and the chamber is evacuated to a vacuum of 28 inches mercury. The tissue is placed approximately 3.5 inches away from the retaining screen and bombarded three times. Following bombardment, the tissue can be divided in half and placed back into liquid and cultured as described above.

10 Five to seven days post bombardment, the liquid media may be exchanged with fresh media, and eleven to twelve days post bombardment with fresh media containing 50 mg/mL hygromycin. This selective media can be refreshed weekly. Seven to eight weeks post bombardment, green, transformed tissue may be observed growing from untransformed, necrotic embryogenic clusters. Isolated green tissue is removed and inoculated into  
15 individual flasks to generate new, clonally propagated, transformed embryogenic suspension cultures. Each new line may be treated as an independent transformation event. These suspensions can then be subcultured and maintained as clusters of immature embryos or regenerated into whole plants by maturation and germination of individual somatic embryos.

#### EXAMPLE 6

##### 20 Expression of Chimeric Genes in Microbial Cells

The cDNAs encoding the instant polypeptides can be inserted into the T7 *E. coli* expression vector pBT430. This vector is a derivative of pET-3a (Rosenberg et al. (1987) *Gene* 56:125-135) which employs the bacteriophage T7 RNA polymerase/T7 promoter system. Plasmid pBT430 was constructed by first destroying the EcoR I and Hind III sites in  
25 pET-3a at their original positions. An oligonucleotide adaptor containing EcoR I and Hind III sites was inserted at the BamH I site of pET-3a. This created pET-3aM with additional unique cloning sites for insertion of genes into the expression vector. Then, the Nde I site at the position of translation initiation was converted to an Nco I site using oligonucleotide-directed mutagenesis. The DNA sequence of pET-3aM in this region,  
30 5'-CATATGG, was converted to 5'-CCCATGG in pBT430.

Plasmid DNA containing a cDNA may be appropriately digested to release a nucleic acid fragment encoding the protein. This fragment may then be purified on a 1% NuSieve GTG™ low melting agarose gel (FMC). Buffer and agarose contain 10  $\mu$ g/ml ethidium bromide for visualization of the DNA fragment. The fragment can then be purified from the  
35 agarose gel by digestion with GELase™ (Epicentre Technologies) according to the manufacturer's instructions, ethanol precipitated, dried and resuspended in 20  $\mu$ L of water. Appropriate oligonucleotide adapters may be ligated to the fragment using T4 DNA ligase (New England Biolabs, Beverly, MA). The fragment containing the ligated adapters can be

purified from the excess adapters using low melting agarose as described above. The vector pBT430 is digested, dephosphorylated with alkaline phosphatase (NEB) and deproteinized with phenol/chloroform as described above. The prepared vector pBT430 and fragment can then be ligated at 16°C for 15 hours followed by transformation into DH5 electrocompetent cells (GIBCO BRL). Transformants can be selected on agar plates containing LB media and 100 µg/mL ampicillin. Transformants containing the gene encoding the instant polypeptides are then screened for the correct orientation with respect to the T7 promoter by restriction enzyme analysis.

For high level expression, a plasmid clone with the cDNA insert in the correct orientation relative to the T7 promoter can be transformed into *E. coli* strain BL21(DE3) (Studier et al. (1986) *J. Mol. Biol.* 189:113-130). Cultures are grown in LB medium containing ampicillin (100 mg/L) at 25°C. At an optical density at 600 nm of approximately 1, IPTG (isopropylthio-β-galactoside, the inducer) can be added to a final concentration of 0.4 mM and incubation can be continued for 3 h at 25°. Cells are then harvested by centrifugation and re-suspended in 50 µL of 50 mM Tris-HCl at pH 8.0 containing 0.1 mM DTT and 0.2 mM phenyl methylsulfonyl fluoride. A small amount of 1 mm glass beads can be added and the mixture sonicated 3 times for about 5 seconds each time with a microprobe sonicator. The mixture is centrifuged and the protein concentration of the supernatant determined. One µg of protein from the soluble fraction of the culture can be separated by SDS-polyacrylamide gel electrophoresis. Gels can be observed for protein bands migrating at the expected molecular weight.

Various modifications of the invention in addition to those shown and described herein will be apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims.

The disclosure of each reference set forth above is incorporated herein by reference in its entirety.

CLAIMS

What is claimed is:

1. An isolated polynucleotide comprising a first nucleotide sequence encoding a polypeptide of at least 348 amino acids that has at least 80% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of  
5 SEQ ID NOs:2, 4, 6, 8 and 10 or a second nucleotide sequence comprising the complement of the first nucleotide sequence.
2. The isolated polynucleotide of Claim 1, wherein the first nucleotide sequence consists of a nucleic acid sequence selected from the group consisting of SEQ ID NOs:1, 3,  
10 5, 7 and 9 that codes for the polypeptide selected from the group consisting of SEQ ID NOs:2, 4, 6, 8 and 10.
3. The isolated polynucleotide of Claim 1 wherein the nucleotide sequences are DNA.
4. The isolated polynucleotide of Claim 1 wherein the nucleotide sequences are  
15 RNA.
5. A chimeric gene comprising the isolated polynucleotide of Claim 1 operably linked to suitable regulatory sequences.
6. A host cell comprising the chimeric gene of Claim 5.
7. A host cell comprising an isolated polynucleotide of Claim 1.
- 20 8. The host cell of Claim 7 wherein the host cell is selected from the group consisting of yeast, bacteria, plant, and virus.
9. A virus comprising the isolated polynucleotide of Claim 1.
10. A polypeptide of at least 348 amino acids that has at least 80% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group  
25 consisting of SEQ ID NOs:2, 4, 6, 8 and 10.
11. A method of selecting an isolated polynucleotide that affects the level of expression of a flavonoid biosynthetic enzyme polypeptide in a plant cell, the method comprising the steps of:
  - (a) constructing an isolated polynucleotide comprising a nucleotide sequence of  
30 at least one of 30 contiguous nucleotides derived from an isolated polynucleotide of Claim 1;
  - (b) introducing the isolated polynucleotide into a plant cell;
  - (c) measuring the level of a polypeptide in the plant cell containing the polynucleotide; and
  - 35 (d) comparing the level of polypeptide in the plant cell containing the isolated polynucleotide with the level of polypeptide in a plant cell that does not contain the isolated polynucleotide.

12. The method of Claim 11 wherein the isolated polynucleotide consists of a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7 and 9 that codes for the polypeptide selected from the group consisting of SEQ ID NOs:2, 4, 6, 8 and 10.

5 13. A method of selecting an isolated polynucleotide that affects the level of expression of a flavonoid biosynthetic enzyme polypeptide in a plant cell, the method comprising the steps of:

(a) constructing an isolated polynucleotide of Claim 1;

(b) introducing the isolated polynucleotide into a plant cell; and

10 (c) measuring the level of polypeptide in the plant cell containing the polynucleotide to provide a positive selection means.

14. A method of obtaining a nucleic acid fragment encoding a flavonoid biosynthetic enzyme polypeptide comprising the steps of:

15 (a) synthesizing an oligonucleotide primer comprising a nucleotide sequence of at least one of 30 contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9 and the complement of such nucleotide sequences; and

(b) amplifying a nucleic acid sequence using the oligonucleotide primer.

20 15. A method of obtaining a nucleic acid fragment encoding a flavonoid biosynthetic enzyme polypeptide comprising the steps of:

(a) probing a cDNA or genomic library with an isolated polynucleotide comprising at least one of 30 contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9 and the complement of such nucleotide sequences;

25 (b) identifying a DNA clone that hybridizes with the isolated polynucleotide;

(c) isolating the identified DNA clone; and

(d) sequencing the cDNA or genomic fragment that comprises the isolated DNA clone.

16. A composition comprising the isolated polynucleotide of Claim 1.

30 17. A composition comprising the isolated polypeptide of Claim 10.

18. An isolated polynucleotide comprising the nucleotide sequence having at least one of 30 contiguous nucleotides derived from a nucleic acid sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9 and the complement of such sequences.

35 19. An expression cassette comprising an isolated polynucleotide of Claim 1 operably linked to a promoter.

20. A method for positive selection of a transformed cell comprising:

(a) transforming a host cell with the chimeric gene of Claim 5 or an expression cassette of Claim 19; and

(b) growing the transformed host cell under conditions which allow expression of the polynucleotide in an amount sufficient to alter isoflavone levels in the cell to provide a positive selection means.

21. The method of Claim 20 wherein the host cell is a plant cell.
- 5     22. The method of any of Claims 11 or 13 wherein the plant cell is a monocot.
23. The method of any of Claims 11 or 13 wherein the plant cell is a dicot.

SEQUENCE LISTING

&lt;110&gt; E. I. du Pont de Nemours and Company

&lt;120&gt; Flavonoid Biosynthetic Enzymes

&lt;130&gt; BB1324 1

&lt;140&gt;

&lt;141&gt;

&lt;150&gt; 60/113,190

&lt;151&gt; 1998-12-21

&lt;160&gt; 10

&lt;170&gt; Microsoft Office 97

&lt;210&gt; 1

&lt;211&gt; 1053

&lt;212&gt; DNA

&lt;213&gt; Glycine max

&lt;400&gt; 1

```

gataacccaaa aagcaatgga actctttgag ggccaatctc ttttgtacat gcagctatac 60
gggcacctaag gacctatgtg tcttaagtgg gctgttcaac taggtattcc agacataata 120
cagaaccatg ccaaaccat tacagtttct gacttggttt ctactcttca aatttcacca 180
tctaaggctg gttttgtgca gcagttcatg cgctttttgg cacacgatgg aatctttgat 240
atcogtgaga gccaagatga tcatgaatta gcatatgctc taaccctgc atcaaagctt 300
ctagttagtt gcagtgacca ctgtttatct ccaatgggtc ggatgaatac tgatccactt 360
ctgatgacta cataccatca ctttggggaa tggattcgtg gggaagacc cagctacat 420
gagacagcct tcggaacaag ctttggggga cttcttgaga aaaacctac acaaagtgt 480
ctcttcaatg aggctatggc aagtgttcc cgaatggtag acttggcact caaaaattgc 540
acttcagttt ttgaagggt agattccatg gtggatgtt gtggtggaac tgggaaccaca 600
gccaaaatta tctgtgaggc atttccgaag ttgaaatgtg ttgtgcttga ccttcctcat 660
gtttagagaaa acttgacagg aaccaataat ttgagttttg tcggtggtga tatgttcaac 720
tctttccctc aaactgatgc agttctacta aagtgggttt tacataattg gaatgacgaa 780
aattgcataa agatcctgaa aaagtgtaaa gattctattt caagcaaagg caacaaagga 840
aaagtgatta tcatagatat aataataaat gagaagctag atgatccgga tatgactcga 900
acaaagctta gtttgatgat agttatgtcg actatgaatg gaagagagcg aagtgaaaaa 960
gaatggaaaac aaatgttcat tgaagcggga ttcaaact gcaaaatatt tcccatcttt 1020
ggtttcagat ctctaattga gctctatcct tag 1053

```

&lt;210&gt; 2

&lt;211&gt; 350

&lt;212&gt; PRT

&lt;213&gt; Glycine max

&lt;400&gt; 2

```

Asp Asn Gln Lys Ala Met Glu Leu Phe Glu Gly Gln Ser Leu Leu Tyr
  1              5              10              15

```

```

Met Gln Leu Tyr Gly His Leu Arg Pro Met Cys Leu Lys Trp Ala Val
      20              25              30

```

```

Gln Leu Gly Ile Pro Asp Ile Ile Gln Asn His Ala Lys Pro Ile Thr
      35              40              45

```

```

Val Ser Asp Leu Val Ser Thr Leu Gln Ile Ser Pro Ser Lys Ala Gly
      50              55              60

```

Phe Val Gln Gln Phe Met Arg Phe Leu Ala His Asp Gly Ile Phe Asp  
 65 70 75 80  
 Ile Arg Glu Ser Gln Asp Asp His Glu Leu Ala Tyr Ala Leu Thr Pro  
 85 90 95  
 Ala Ser Lys Leu Leu Val Ser Cys Ser Asp His Cys Leu Ser Pro Met  
 100 105 110  
 Val Arg Met Asn Thr Asp Pro Leu Leu Met Thr Thr Tyr His His Phe  
 115 120 125  
 Gly Glu Trp Ile Arg Gly Glu Asp Pro Thr Val His Glu Thr Ala Phe  
 130 135 140  
 Gly Thr Ser Phe Trp Gly Leu Leu Glu Lys Asn Pro Thr Gln Met Ser  
 145 150 155 160  
 Leu Phe Asn Glu Ala Met Ala Ser Asp Ser Arg Met Val Asp Leu Ala  
 165 170 175  
 Leu Lys Asn Cys Thr Ser Val Phe Glu Gly Leu Asp Ser Met Val Asp  
 180 185 190  
 Val Gly Gly Gly Thr Gly Thr Thr Ala Lys Ile Ile Cys Glu Ala Phe  
 195 200 205  
 Pro Lys Leu Lys Cys Val Val Leu Asp Leu Pro His Val Val Glu Asn  
 210 215 220  
 Leu Thr Gly Thr Asn Asn Leu Ser Phe Val Gly Gly Asp Met Phe Asn  
 225 230 235 240  
 Ser Phe Pro Gln Thr Asp Ala Val Leu Leu Lys Trp Val Leu His Asn  
 245 250 255  
 Trp Asn Asp Glu Asn Cys Ile Lys Ile Leu Lys Lys Cys Lys Asp Ser  
 260 265 270  
 Ile Ser Ser Lys Gly Asn Lys Gly Lys Val Ile Ile Ile Asp Ile Ile  
 275 280 285  
 Ile Asn Glu Lys Leu Asp Asp Pro Asp Met Thr Arg Thr Lys Leu Ser  
 290 295 300  
 Leu Asp Ile Val Met Ser Thr Met Asn Gly Arg Glu Arg Ser Glu Lys  
 305 310 315 320  
 Glu Trp Lys Gln Met Phe Ile Glu Ala Gly Phe Lys His Cys Lys Ile  
 325 330 335  
 Phe Pro Ile Phe Gly Phe Arg Ser Leu Ile Glu Leu Tyr Pro  
 340 345 350

&lt;210&gt; 3

&lt;211&gt; 1160

&lt;212&gt; DNA

&lt;213&gt; Glycine max

<400> 3  
 gtttgctatt atggcttctt cattaaacaa tggccgtaaa gcaagtgaga tttttcaagg 60  
 tcaagctctc ttgtacaaac atttgcttgg cttcatagat tctaagtgtc taaaatggat 120  
 gggtgagctt gacatacccg acataatcca cagccatagc catggccaac ccattacttt 180  
 ttcagagttg gtgtcaattc tacaagtccc accaactaaa actcgtcagg tgcagagcct 240  
 catgcgttat ctagcacaca atggattctt tgagatagta agaatccatg acaacataga 300  
 agcatatgct ctactgctg cttcagagtt acttgtcaaa agcagtgagc ttagtttagc 360  
 tccaatgggt gagtattttc ttgaaccaa ttgtcaaggt gcatggaacc agttgaagag 420  
 gtgggttcat gaggaagatc tcacagtatt tgaggtctcc ttaggaacac ctttctggga 480  
 ctttatcaat aaagaccctg catataacaa gtcattcaat gaggcaatgg cttgtgattc 540  
 tcagatgttg aacttgccgt ttagagattg caattgggtc tttgaggac tggaatccat 600  
 tgtggatgtt ggtggtggaa ctggaatcac agcaaagatt atctgtgagg cttttcctaa 660  
 gctgaaatgc atggtgttg aacgtccaaa tgttgtggaa aatttgtcag gaagcaacaa 720  
 tttgacattt gttggtgggg acatgtttta atgcatcccc aaggctgatg cagttctgct 780  
 taagttggtt ttacataatt ggaatgacaa cgattgcatg aagatattag aaaattgtaa 840  
 agaagctatt tcaggtgaaa gcaaaacagg aaaagtagtt gtcatagata ctgtgataaa 900  
 cgaaaacaaa gatgagcgcc aagttactga actaaagctc cttatggatg tacacatggc 960  
 atgtattatt aatggaaaag agagaaaaga agaagattgg aagaaactct tcatggaagc 1020  
 aggggttccaa agctacaaaa tatctccctt cacaggatat ttgtctctta ttgagatcta 1080  
 tccttgaata ctgacgtgc aatattccat ttagtagtta atttgcattg tatcaataaa 1140  
 ataaaagtta tctttgtgtt 1160

<210> 4  
 <211> 358  
 <212> PRT  
 <213> Glycine max

<400> 4  
 Met Ala Ser Ser Leu Asn Asn Gly Arg Lys Ala Ser Glu Ile Phe Gln  
 1 5 10 15  
 Gly Gln Ala Leu Leu Tyr Lys His Leu Leu Gly Phe Ile Asp Ser Lys  
 20 25 30  
 Cys Leu Lys Trp Met Val Glu Leu Asp Ile Pro Asp Ile Ile His Ser  
 35 40 45  
 His Ser His Gly Gln Pro Ile Thr Phe Ser Glu Leu Val Ser Ile Leu  
 50 55 60  
 Gln Val Pro Pro Thr Lys Thr Arg Gln Val Gln Ser Leu Met Arg Tyr  
 65 70 75 80  
 Leu Ala His Asn Gly Phe Phe Glu Ile Val Arg Ile His Asp Asn Ile  
 85 90 95  
 Glu Ala Tyr Ala Leu Thr Ala Ala Ser Glu Leu Leu Val Lys Ser Ser  
 100 105 110  
 Glu Leu Ser Leu Ala Pro Met Val Glu Tyr Phe Leu Glu Pro Asn Cys  
 115 120 125  
 Gln Gly Ala Trp Asn Gln Leu Lys Arg Trp Val His Glu Glu Asp Leu  
 130 135 140  
 Thr Val Phe Glu Val Ser Leu Gly Thr Pro Phe Trp Asp Phe Ile Asn  
 145 150 155 160  
 Lys Asp Pro Ala Tyr Asn Lys Ser Phe Asn Glu Ala Met Ala Cys Asp  
 165 170 175



Ser Gln Met Leu Asn Leu Ala Phe Arg Asp Cys Asn Trp Val Phe Glu  
 180 185 190

Gly Leu Glu Ser Ile Val Asp Val Gly Gly Gly Thr Gly Ile Thr Ala  
 195 200 205

Lys Ile Ile Cys Glu Ala Phe Pro Lys Leu Lys Cys Met Val Leu Glu  
 210 215 220

Arg Pro Asn Val Val Glu Asn Leu Ser Gly Ser Asn Asn Leu Thr Phe  
 225 230 235 240

Val Gly Gly Asp Met Phe Lys Cys Ile Pro Lys Ala Asp Ala Val Leu  
 245 250 255

Leu Lys Leu Val Leu His Asn Trp Asn Asp Asn Asp Cys Met Lys Ile  
 260 265 270

Leu Glu Asn Cys Lys Glu Ala Ile Ser Gly Glu Ser Lys Thr Gly Lys  
 275 280 285

Val Val Val Ile Asp Thr Val Ile Asn Glu Asn Lys Asp Glu Arg Gln  
 290 295 300

Val Thr Glu Leu Lys Leu Leu Met Asp Val His Met Ala Cys Ile Ile  
 305 310 315 320

Asn Gly Lys Glu Arg Lys Glu Glu Asp Trp Lys Lys Leu Phe Met Glu  
 325 330 335

Ala Gly Phe Gln Ser Tyr Lys Ile Ser Pro Phe Thr Gly Tyr Leu Ser  
 340 345 350

Leu Ile Glu Ile Tyr Pro  
 355

<210> 5  
 <211> 1065  
 <212> DNA  
 <213> Glycine max

<400> 5

atggcttcaa	tgaataacca	aaaagaaatt	gagctctttg	agggccaatc	tcttctgtac	60
atgcagctat	atgggcacct	aagacctatg	tgtcttaagt	gggctgttca	actaggtatt	120
ccagacataa	tacagaacca	tgccaaaccc	atttctcttt	ctgacttggt	ctctactctt	180
caaattccac	cagctaacgc	tgcttttggt	cagcggttca	tgcgcttctt	ggcacacaat	240
ggaatctttg	agatccatga	gagccaagaa	gatcatgaac	taacatatgc	tctaaccctt	300
gcatcaaagc	ttcttggtcaa	tagtagtgat	cattgtctat	ctccaatggt	tctagcgttt	360
accgatccac	ttcggaacgt	taaataccat	cacttggggg	aatggattcg	tggggaggac	420
ccctcagtat	ttgagacagc	ccacggaaca	agcgcttggg	gacttcttga	gaaaaatcct	480
gaatatttta	gtctcttcaa	tgaggctatg	gcaagtgatt	cccgaatagt	agacttggca	540
ctcaaaaatt	gcacttcagt	ttttgagggg	ctagattcca	tggtggatgt	tggtgggtgga	600
actggaacca	cggccagaat	tatctgtgac	gcatttccta	agttgaaatg	tggtgtgctt	660
gaccttcctc	atgttgtaga	gaacttgaca	gggaccaata	atttgagttt	tggtgggtggt	720
gacatgttca	actctatccc	tcaagctgat	gcagtgctac	taaagtgggt	tttacataat	780
tggaaccgacg	aaaattgcat	aaagatcctg	caaaagtgtg	gagattctat	ttcaagcaaa	840
ggcaacagtg	gaaaagtgat	tatcatagat	gccgtaataa	atgagaagct	agatgaccog	900
gatatgacac	aaacaaagct	tagtttggtg	attattatgt	tgacgatgaa	tggaagagag	960

agaacggaaa aagaatggaa acaactcttc atcgaagcag gattcaaaca ctacaaaata 1020  
 tttcccatct ttggttttag atctctgatt gaggtctatc cttga 1065

<210> 6  
 <211> 351  
 <212> PRT  
 <213> Glycine max

<400> 6  
 Met Ala Ser Met Asn Asn Gln Lys Glu Ile Glu Leu Phe Glu Gly Gln  
 1 5 10 15  
 Ser Leu Leu Tyr Met Gln Leu Tyr Gly His Leu Arg Pro Met Cys Leu  
 20 25 30  
 Lys Trp Ala Val Gln Leu Gly Ile Pro Asp Ile Ile Gln Asn His Ala  
 35 40 45  
 Lys Pro Ile Ser Leu Ser Asp Leu Val Ser Thr Leu Gln Ile Pro Pro  
 50 55 60  
 Ala Asn Ala Ala Phe Val Gln Arg Phe Met Arg Phe Leu Ala His Asn  
 65 70 75 80  
 Gly Ile Phe Glu Ile His Glu Ser Gln Glu Leu Thr Tyr Ala Leu Thr  
 85 90 95  
 Pro Ala Ser Lys Leu Leu Val Asn Ser Ser Asp His Cys Leu Ser Pro  
 100 105 110  
 Met Val Leu Ala Phe Thr Asp Pro Leu Arg Asn Val Lys Tyr His His  
 115 120 125  
 Leu Gly Glu Trp Ile Arg Gly Glu Asp Pro Ser Val Phe Glu Thr Ala  
 130 135 140  
 His Gly Thr Ser Ala Trp Gly Leu Leu Glu Lys Asn Pro Glu Tyr Phe  
 145 150 155 160  
 Ser Leu Phe Asn Glu Ala Met Ala Ser Asp Ser Arg Ile Val Asp Leu  
 165 170 175  
 Ala Leu Lys Asn Cys Thr Ser Val Phe Glu Gly Leu Asp Ser Met Val  
 180 185 190  
 Asp Val Gly Gly Gly Thr Gly Thr Thr Ala Arg Ile Ile Cys Asp Ala  
 195 200 205  
 Phe Pro Lys Leu Lys Cys Val Val Leu Asp Leu Pro His Val Val Glu  
 210 215 220  
 Asn Leu Thr Gly Thr Asn Asn Leu Ser Phe Val Gly Gly Asp Met Phe  
 225 230 235 240  
 Asn Ser Ile Pro Gln Ala Asp Ala Val Leu Leu Lys Trp Val Leu His  
 245 250 255  
 Asn Trp Thr Asp Glu Asn Cys Ile Lys Ile Leu Gln Lys Cys Arg Asp  
 260 265 270

Ser Ile Ser Ser Lys Gly Asn Ser Gly Lys Val Ile Ile Ile Asp Ala  
275 280 285

Val Ile Asn Glu Lys Leu Asp Asp Pro Asp Met Thr Gln Thr Lys Leu  
290 295 300

Ser Leu Asp Ile Ile Met Leu Thr Met Asn Gly Arg Glu Arg Thr Glu  
305 310 315 320

Lys Glu Trp Lys Gln Leu Phe Ile Glu Ala Gly Phe Lys His Tyr Lys  
325 330 335

Ile Phe Pro Ile Phe Gly Phe Arg Ser Leu Ile Glu Val Tyr Pro  
340 345 350

<210> 7  
<211> 1253  
<212> DNA  
<213> Glycine max

<400> 7  
ccaaaaagca atggaactct ttgagggcca atctcttttg tacatgcagc tatatgggca 60  
cctaagacct atgtgtctta agtgggctgt tcaactaggt attccagaca taatacagaa 120  
ccatgccaaa cccattactc tttctgattt ggtctctact cttcaaattc caccatctaa 180  
ggctgggttt gtgcagcagt tcatgcgctt tttggcacac gatggaatct ttgatatccg 240  
tgagagccaa gatgatcatg aattagcata tgctctaacc cctgcttcaa agcttctagt 300  
tagttgcagt gaccactgtt tatctccaat gggtcggatg aatactgatc cacttctgat 360  
gactacatac catcactttg ggggaatggat tcgtggggaa gacccacag tacatgagac 420  
agccttcgga acaagctttt ggggacttct tgagaaaaaac cctacacaaa tgagtctctt 480  
caatgaggct atggcaagtg attcccgaat ggtagacttg gcactcaaaa attgcacttc 540  
agtttttgaa gggctagatt ccatgggtgga tgttggtggt ggaactggaa ccacagccag 600  
aattatctgt gaggcatttc cgaagttaga atgtgttggt cttgaccttc ctcatgttgt 660  
agaaaacttg acaggaacca ataatttgag tttgtcggt ggtgacatgt tcaactcttt 720  
ccctcaaact gatgcagttc tactaaagtg ggttttacat aattggaatg acgaaaattg 780  
cataaagatc ctgaaaaagt gtaaagattc tatttcaagc aaaggcaaca gaggaaaagt 840  
gattatcata gatataataa taaatgagaa gctagatgac ccggatatga ctgaaacaaa 900  
gcttagtttg gatatagtta tgtcgactat gaatggaaga gagcgaagtg aaaaagaatg 960  
gaaacaaatg ttcattgaag cgggattcca acactgcaaa atatttccca tctttggttt 1020  
cagatctcta attgagctct atccttagac atatactcta gcttgttatg cattttatag 1080  
tttgcgcatg ttgtgatgag ctgaataaag ctatgatggc atagcattgt caacttctgg 1140  
tggaattaca atatgttcc taccactagc tatagggtgt taaatgttca gtgtacggat 1200  
ttctcattcc tatttggccaa aaataaattg aatgaacatg taatggattt agt 1253

<210> 8  
<211> 348  
<212> PRT  
<213> Glycine max

<400> 8  
Gln Lys Ala Met Glu Leu Phe Glu Gly Gln Ser Leu Leu Tyr Met Gln  
1 5 10 15  
Leu Tyr Gly His Leu Arg Pro Met Cys Leu Lys Trp Ala Val Gln Leu  
20 25 30  
Gly Ile Pro Asp Ile Ile Gln Asn His Ala Lys Pro Ile Thr Leu Ser  
35 40 45  
Asp Leu Val Ser Thr Leu Gln Ile Pro Pro Ser Lys Ala Gly Phe Val  
50 55 60

Gln Gln Phe Met Arg Phe Leu Ala His Asp Gly Ile Phe Asp Ile Arg  
 65 70 75 80  
 Glu Ser Gln Asp Asp His Glu Leu Ala Tyr Ala Leu Thr Pro Ala Ser  
 85 90 95  
 Lys Leu Leu Val Ser Cys Ser Asp His Cys Leu Ser Pro Met Val Arg  
 100 105 110  
 Met Asn Thr Asp Pro Leu Leu Met Thr Thr Tyr His His Phe Gly Glu  
 115 120 125  
 Trp Ile Arg Gly Glu Asp Pro Thr Val His Glu Thr Ala Phe Gly Thr  
 130 135 140  
 Ser Phe Trp Gly Leu Leu Glu Lys Asn Pro Thr Gln Met Ser Leu Phe  
 145 150 155 160  
 Asn Glu Ala Met Ala Ser Asp Ser Arg Met Val Asp Leu Ala Leu Lys  
 165 170 175  
 Asn Cys Thr Ser Val Phe Glu Gly Leu Asp Ser Met Val Asp Val Gly  
 180 185 190  
 Gly Gly Thr Gly Thr Thr Ala Arg Ile Ile Cys Glu Ala Phe Pro Lys  
 195 200 205  
 Leu Lys Cys Val Val Leu Asp Leu Pro His Val Val Glu Asn Leu Thr  
 210 215 220  
 Gly Thr Asn Asn Leu Ser Phe Val Gly Gly Asp Met Phe Asn Ser Phe  
 225 230 235 240  
 Pro Gln Thr Asp Ala Val Leu Leu Lys Trp Val Leu His Asn Trp Asn  
 245 250 255  
 Asp Glu Asn Cys Ile Lys Ile Leu Lys Lys Cys Lys Asp Ser Ile Ser  
 260 265 270  
 Ser Lys Gly Asn Arg Gly Lys Val Ile Ile Ile Asp Ile Ile Ile Asn  
 275 280 285  
 Glu Lys Leu Asp Asp Pro Asp Met Thr Arg Thr Lys Leu Ser Leu Asp  
 290 295 300  
 Ile Val Met Ser Thr Met Asn Gly Arg Glu Arg Ser Glu Lys Glu Trp  
 305 310 315 320  
 Lys Gln Met Phe Ile Glu Ala Gly Phe Gln His Cys Lys Ile Phe Pro  
 325 330 335  
 Ile Phe Gly Phe Arg Ser Leu Ile Glu Leu Tyr Pro  
 340 345

&lt;210&gt; 9

&lt;211&gt; 1065

&lt;212&gt; DNA

&lt;213&gt; Glycine max

<400> 9  
 atggcttcaa tgaataacca aaaagaaatt gagctctttg agggccaatc tcttctgtac 60  
 atgcagctat atgggcacct aagacctatg tgtcttaagt gggctgttca actaggtatt 120  
 ccagacataa tacagaacca tgccaaaccc atttctcttt ctgacttggt ctctactctt 180  
 caaattccac cagctaacgc tgctttttgtg cagcggttca tgcgcttctt ggcacacaat 240  
 ggaatctttg agatccatga gagccaagaa gatcatgaac taacatatgc tctaaccctt 300  
 gcatcaaagc ttcttgtcaa tagtagtgat cattgtctat ctccaatggt tctagcgttt 360  
 accgatccac ttcggaacgt taaataccat cacttggggg aatggattcg tggggaggac 420  
 ccctcagtat ttgagacagc ccacggaaca agcgcttggg gacttcttga gaaaaatcct 480  
 gaatatctta gtctcttcaa tgaggctatg gcaagtgatt cccgaatagt agacttggca 540  
 ctcaaaaatt gcacttcagt ttttgagggg ctagattcca tgggtgatgt tgggtggtgga 600  
 actggaacca cggccagaat tatctgtgac gcatttccta agttgaaatg tgttgtgctt 660  
 gaccttcctc atgttgtaga gaacttgaca gggaccaata atttgagttt tgttgggtggt 720  
 gacatgttca actctatccc tcaagctgat gcagtgtac taaagtgggt ttacataat 780  
 tggaccgacg aaaattgcat aaagatcctg caaaagtgt gagattctat ttcaagcaaa 840  
 ggcaacagtg gaaaagtgat tatcatagat gccgtaataa atgagaagct agatgacctg 900  
 gatatgacac aaacaaagct tagtttgac attattatgt tgacgatgaa tggagagag 960  
 agaacggaaa aagaatggaa acaactcttc atcgaagcag gattcaaaca ctacaaaata 1020  
 tttcccatct ttggttttag atctctgatt gaggtctatc cttga 1065

<210> 10  
 <211> 350  
 <212> PRT  
 <213> Glycine max

<400> 10  
 Ala Ser Met Asn Asn Gln Lys Glu Ile Glu Leu Phe Glu Gly Gln Ser  
 1 5 10 15  
 Leu Leu Tyr Met Gln Leu Tyr Gly His Leu Arg Pro Met Cys Leu Lys  
 20 25 30  
 Trp Ala Val Gln Leu Gly Ile Pro Asp Ile Ile Gln Asn His Ala Lys  
 35 40 45  
 Pro Ile Ser Leu Ser Asp Leu Val Ser Thr Leu Gln Ile Pro Pro Ala  
 50 55 60  
 Asn Ala Ala Phe Val Gln Arg Phe Met Arg Phe Leu Ala His Asn Gly  
 65 70 75 80  
 Ile Phe Glu Ile His Glu Ser Gln Glu Leu Thr Tyr Ala Leu Thr Pro  
 85 90 95  
 Ala Ser Lys Leu Leu Val Asn Ser Ser Asp His Cys Leu Ser Pro Met  
 100 105 110  
 Val Leu Ala Phe Thr Asp Pro Leu Arg Asn Val Lys Tyr His His Leu  
 115 120 125  
 Gly Glu Trp Ile Arg Gly Glu Asp Pro Ser Val Phe Glu Thr Ala His  
 130 135 140  
 Gly Thr Ser Ala Trp Gly Leu Leu Glu Lys Asn Pro Glu Tyr Phe Ser  
 145 150 155 160  
 Leu Phe Asn Glu Ala Met Ala Ser Asp Ser Arg Ile Val Asp Leu Ala  
 165 170 175

Leu Lys Asn Cys Thr Ser Val Phe Glu Gly Leu Asp Ser Met Val Asp  
180 185 190

Val Gly Gly Gly Thr Gly Thr Thr Ala Arg Ile Ile Cys Asp Ala Phe  
195 200 205

Pro Lys Leu Lys Cys Val Val Leu Asp Leu Pro His Val Val Glu Asn  
210 215 220

Leu Thr Gly Thr Asn Asn Leu Ser Phe Val Gly Gly Asp Met Phe Asn  
225 230 235 240

Ser Ile Pro Gln Ala Asp Ala Val Leu Leu Lys Trp Val Leu His Asn  
245 250 255

Trp Thr Asp Glu Asn Cys Ile Lys Ile Leu Gln Lys Cys Arg Asp Ser  
260 265 270

Ile Ser Ser Lys Gly Asn Ser Gly Lys Val Ile Ile Ile Asp Ala Val  
275 280 285

Ile Asn Glu Lys Leu Asp Asp Pro Asp Met Thr Gln Thr Lys Leu Ser  
290 295 300

Leu Asp Ile Ile Met Leu Thr Met Asn Gly Arg Glu Arg Thr Glu Lys  
305 310 315 320

Glu Trp Lys Gln Leu Phe Ile Glu Ala Gly Phe Lys His Tyr Lys Ile  
325 330 335

Phe Pro Ile Phe Gly Phe Arg Ser Leu Ile Glu Val Tyr Pro  
340 345 350

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C12N 15/54, 15/82, 9/10, C12Q 1/02, 1/68, 1/48</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 00/37656</b> <b>(43) International Publication Date:</b> 29 June 2000 (29.06.00)
<b>(21) International Application Number:</b> PCT/US99/30338 <b>(22) International Filing Date:</b> 20 December 1999 (20.12.99)  <b>(30) Priority Data:</b> 60/113,190      21 December 1998 (21.12.98)      US  <b>(71) Applicant (for all designated States except US):</b> E.I. DU PONT DE NEMOURS AND COMPANY [US/US]; 1007 Market Street, Wilmington, DE 19898 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> FAMODU, Omolayo, O. [US/US]; 216 Barrett Run Place, Newark, DE 19702 (US). MCGONIGLE, Brian [US/US]; 1707 North Union Street, Wilmington, DE 19806 (US). ODELL, Joan, T. [US/US]; P.O. Box 826, Unionville, PA 19375 (US). FADER, Gary, M. [US/US]; 1000 Woods Lane, Landenberg, PA 19350 (US). FALCO, Saverio, Carl [US/US]; 1902 Miller Road, Arden, DE 19810 (US).  <b>(74) Agent:</b> FEULNER, Gregory, J.; E.I. du Pont de Nemours and Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).		<b>(81) Designated States:</b> AE, AL, AU, BA, BB, BG, BR, CA, CN, CR, CU, CZ, DM, EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>  <b>(88) Date of publication of the international search report:</b> 14 September 2000 (14.09.00)
<b>(54) Title:</b> FLAVONOID BIOSYNTHETIC ENZYMES  <b>(57) Abstract</b>  This invention relates to an isolated nucleic acid fragment encoding a flavonoid biosynthetic enzyme. The invention also relates to the construction of a chimeric gene encoding all or a portion of the flavonoid biosynthetic enzyme, in sense or antisense orientation, wherein expression of the chimeric gene results in production of altered levels of the flavonoid biosynthetic enzyme in a transformed host cell.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 99/30338

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/54 C12N15/82 C12N9/10 C1201/02 C1201/68  
C1201/48

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	SHOEMAKER R., ET AL.: "sa87h11.y1 Gm-c1004 Glycine max cDNA clone GENOME SYSTEMS CLONE ID:Gm-c1004-6334 5' similar to TR:022308 022308 7-O-METHYLTRANSFERASE.'3! TR:022309 TR:022555 ;, mRNA sequence." EMBL ACCESSION NO:AI507849, 17 March 1999 (1999-03-17), XP002139373 the whole document	1-4, 10, 16-18
P, X	SHOEMAKER, R., ET AL.: "sf38a06.y1 Gm-c1028 Glycine max cDNA clone GENOME SYSTEMS CLONE ID:Gm-c1028-2219 5' similar to TR:022555 022555 O-METHYTRANSFERASE.'2! TR:022308 ;, mRNA sequence." EMBL ACCESSION NO:AW203767, 10 December 1999 (1999-12-10), XP002139374 the whole document	1-4, 10, 16-18



Further documents are listed in the continuation of box C



Patent family members are listed in annex.

### \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "3" document member of the same patent family

Date of the actual completion of the international search

5 June 2000

Date of mailing of the international search report

26/06/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel (+31-70) 340-2040, Tx 31 651 epo nl.  
Fax (+31-70) 340-3016

Authorized officer

Maddox, A

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/30338

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	<p>SHOEMAKER, R., ET AL.: "sa88b04.y1 Gm-cl004 Glycine max cDNA clone GENOME SYSTEMS CLONE ID:Gm-cl004-6344 5' similar to TR:022308 022308 7-0-METHYLTRANSFERASE.'3! TR:022309 TR:022555 ;, mRNA sequence." EMBL ACCESSION NO:AI507864, 17 March 1999 (1999-03-17), XP002139375 the whole document</p>	1-4, 10, 16-18
P, X	<p>SHOEMAKER, R., ET AL.: "sa82g01.y1 Gm-cl004 Glycine max cDNA clone GENOME SYSTEMS CLONE ID:Gm-cl004-5833 5' similar to SW:ZRP4_MAIZE P479170-METHYLTRANSFERASE ZRP4 ;, mRNA sequence." EMBL ACCESSION NO:AI441988, 16 March 1999 (1999-03-16), XP002139376 the whole document</p>	1-4, 10, 16-18
A	<p>HE XIAN-ZHI ET AL: "Stress responses in alfalfa (Medicago sativa L). XXII. cDNA cloning and characterization of an elicitor-inducible isoflavone 7-0-methyltransferase." PLANT MOLECULAR BIOLOGY JAN. 1, 1998, vol. 36, no. 1, 1 January 1998 (1998-01-01), pages 43-54, XP002139377 ISSN: 0167-4412 the whole document -&amp; HE, X.Z., ET AL.: "Medicago sativa 7-0-methyltransferase (7-IOMT(6)) mRNA, complete cds." EMBL ACCESSION NO:AF000975, 5 November 1997 (1997-11-05), XP002139378 -&amp; HE X.Z., ET AL.: "Medicago sativa 7-0-methyltransferase (7-IOMT(9)) mRNA, complete cds." EMBL ACCESSION NO:AF000976, 5 November 1997 (1997-11-05), XP002139379 -&amp; HE X.Z., ET AL.: "Medicago sativa isoflavone-0-methyltransferase mRNA, complete cds." EMBL ACCESSION NO:U97125, 5 November 1997 (1997-11-05), XP002139380</p>	1-23

-/--

# INTERNATIONAL SEARCH REPORT

Patent Application No  
PCT/US 99/30338

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>DATABASE BIOSIS 'Online!  BIOSCIENCES INFORMATION SERVICE,  PHILADELPHIA, PA, US1977  POULTON J E ET AL: "O METHYLATION OF  FLAVONOID SUBSTRATES BY A PARTIALLY  PURIFIED ENZYME FROM SOYBEAN CELL  SUSPENSION CULTURES"  Database accession no. PREV197764047255  XP002139381  abstract  &amp; ARCHIVES OF BIOCHEMISTRY AND BIOPHYSICS  1977,  vol. 180, no. 2, 1977, pages 543-549,  ISSN: 0003-9861</p> <p>-----</p>	1-10

